

Diabetes and Chronic Kidney Disease



NKF National Kidney
Foundation®

Making Lives Better



What is diabetes?

Diabetes mellitus, usually called diabetes or “sugar,” is a condition that occurs when your body does not make enough **insulin** or when your body cannot use normal amounts of insulin properly. Insulin is a hormone that regulates the amount of sugar in your blood. A high blood sugar level can cause problems in many parts of your body.

Are there different types of diabetes?

Yes. The most common ones are **Type 1** and **Type 2**. Type 1 diabetes is responsible for about 10 percent of the cases. It usually begins in childhood and is also called insulin-dependent diabetes mellitus. In this type, your pancreas does not make enough insulin and you have to take insulin injections.

Type 2 diabetes (the most common type of diabetes) usually occurs in people over 45 but is becoming more common in younger people. It is also called non-insulin-dependent diabetes mellitus. In this type, your pancreas makes insulin,

but your body does not use it properly. The high blood sugar level often can be controlled by weight loss, exercise and pills, but insulin may also be needed. Type 2 diabetes is particularly common among African Americans, Hispanic Americans, Asian Americans and American Indians.

How does diabetes affect your body?

Diabetes can cause damage to many parts of your body, especially your kidneys, heart, eyes and nerves. High blood pressure and hardening of the arteries (arteriosclerosis) can develop, which can lead to heart disease.

What does diabetes do to the kidneys?

Diabetes may damage the blood vessels in your kidneys. The first sign of kidney damage is the presence of albumin (a type of protein) in the urine. A sensitive urine test for a tiny amount of albumin (microalbuminuria) helps to detect this damage at an early stage in people with diabetes. Later, kidney function may decrease. Your kidney function is checked by estimating your glomerular filtration rate (GFR) from the results of your blood creatinine test. **When your kidneys are damaged, they cannot clean your blood properly,** and waste products build up in your blood. Your body will retain more water and salt than it should, which can result in weight gain and ankle swelling.

Diabetes also may cause damage to nerves (neuropathy) in your body. This can cause difficulty in emptying your bladder. The pressure that results from your full bladder can back up and injure the kidneys. Also, if urine stays in your bladder for a long time, you may get a urinary tract infection. This is because bacteria grow rapidly in urine with a high sugar level.

How many diabetic patients will get chronic kidney disease?

About a third of patients with diabetes may eventually have kidney disease. Certain groups, such as African Americans, Asian Americans, Hispanic Americans and American Indians may have a higher risk of getting this complication.

What can patients with diabetes do to prevent kidney disease?

Keeping good control of your blood sugar can lower your risk of getting chronic kidney disease. You should have a urine test for albumin (a type of protein) at least once a year. You should have your blood pressure checked as often as your doctor recommends, and take blood pressure medicine if your doctor orders it for you. You should have blood tests to check your blood sugar control and to check your kidney function from your blood creatinine level, follow your diabetic diet and get regular exercise. Avoid alcohol and cigarettes. See your doctor as often as you are told.

Many people with diabetes do not get kidney disease. Having diabetes does not always mean your kidneys will fail. Talk to your doctor about your chances of getting chronic kidney disease.



What are the early signs of chronic kidney disease in patients with diabetes?

If you have diabetes, the earliest sign of chronic kidney disease is the presence of albumin in the urine. This is present long before there is evidence of kidney disease in the usual blood tests done in your doctor's office. This is not only an early sign of kidney disease, but also of blood vessel abnormalities that may lead to heart disease. Thus, it is important to ask your doctor about having a urine test for a tiny amount of albumin (microalbuminuria) on a yearly basis. You should have a simple blood test for serum creatinine to estimate the filtering ability of your kidneys—called your glomerular filtration rate or GFR.

TABLE 1

Signs of Kidney Disease in Patients with Diabetes

Early Signs:

- ❖ Albumin in the urine (also indicates an increased risk for heart disease)
- ❖ Going to the bathroom more often at night
- ❖ High blood pressure

Late Signs:

- ❖ Ankle and leg swelling, leg cramps
- ❖ High levels of BUN and a decrease in your glomerular filtration rate (GFR)
- ❖ Less need for insulin or anti-diabetic pills
- ❖ Weakness, paleness and anemia
- ❖ Itching
- ❖ Morning sickness, nausea and vomiting



You will use the bathroom more at night. **Your blood pressure may get too high.** As a person with diabetes, you should have your blood, urine and blood pressure checked regularly. This will lead to better control of your disease and early treatment of high blood pressure and kidney disease.

What are the late signs of kidney disease in patients with diabetes?

Late signs may include **weight gain** and ankle **swelling (edema)**. As your kidney disease progresses, you will also have a **rise in your blood urea nitrogen (BUN) and a decrease in your glomerular filtration rate (GFR)**. You may have **nausea, vomiting, a loss of appetite, weakness, increasing tiredness, itching, muscle cramps (especially in your legs) and a low blood count (anemia)**. You may find you need to use less insulin. This is because diseased kidneys cause less breakdown of insulin. If you have any of these signs, call your doctor.

If diabetes has affected the kidneys, what can be done?

If you have any of the signs of kidney disease listed earlier, you need to call your doctor. With blood and urine tests, your doctor can tell how your kidneys are working. This will help your doctor to order the best treatment for you. Early detection of kidney disease, with the right treatment, can prevent it from getting worse.

What will happen if your kidney function is less than normal?

First, the doctor needs to find out if your diabetes has caused the injury. Other diseases can cause kidney injury. If no other problems are found, your doctor will try to keep your kidneys working as long as possible. The use of a special type of high blood pressure medicine called **angiotensin converting enzyme (ACE) inhibitors** has been shown to help slow the loss of kidney function and reduce heart disease in diabetes. A related type of medicine, **angiotensin II receptor blockers (ARBs)**, has also been shown to benefit patients with diabetes and chronic kidney disease. The following things can help your kidneys work better and last longer:

- ❖ Controlling your blood sugar
- ❖ Controlling high blood pressure
- ❖ Restricting the amount of salt in your diet to help control high blood pressure and reduce body swelling
- ❖ Treatment of urinary tract infections
- ❖ Correction of any problems in your urinary system
- ❖ Avoiding any medicines that may damage the kidneys (especially anti-inflammatory pain-relieving medicines)
- ❖ Checking with your doctor before taking any herbal supplements.



How are the kidneys kept working as long as possible?

The kidney doctor (nephrologist) will plan your treatment with you, your family and your dietitian. The best things to keep your kidneys working are controlling your blood sugar and high blood pressure with an ACE inhibitor or ARB. Your blood pressure should be 125/75 or less. In many cases, more than one high blood pressure medicine may be needed to reach this target.

You should also keep your blood sugar well controlled. A test called **hemoglobin A-1-c** is used to check your average blood sugar. Your result on this test should be less than seven percent. Restriction of protein in your diet may be helpful. You and your dietitian will plan your diet together. (See National Kidney Foundation brochure *Nutrition and Chronic Kidney Disease*.)

What about a low-protein diet?

Research suggests that **lowering the protein in your diet** can slow the advance of kidney

damage. You should talk to your doctor about this. If you need to go on a low-protein diet, you need to plan this with a dietitian. Do not go on this type of diet without talking to a dietitian, as it could make you sicker.

What is kidney failure in patients with diabetes?

Kidney failure means that your kidneys are no longer able to support you in a reasonably healthy state and **dialysis** or **transplantation** is needed. This happens when your kidney function is about 15 percent of normal. **Without proper treatment**, the time between the beginning of diabetic kidney damage and end stage kidney failure is about five to seven years.

How is kidney failure treated in diabetic patients?

Three types of treatment can be used once your kidneys have failed: **kidney transplantation**, **hemodialysis** and **peritoneal dialysis**. The type of treatment for you will be chosen according to your general health and medical condition, your lifestyle and your treatment preference. The success rate of each treatment type is very important in this planning. These decisions are not final. Many people have used each one of these treatments at different times. Your health care team will discuss these different treatments with you and answer your questions.

Can a patient with diabetes have a kidney transplant?

Yes, a kidney transplant can come from someone who has died or from a living donor who may be a close relative, friend or even a stranger who wished to donate a kidney to anyone in need of a transplant. Once you get a new kidney, you may



need to use a higher dose of insulin. This is because you will be eating more, your new kidney will break down insulin better than your injured one, and you will be using medicines called steroids to keep your body from rejecting your new kidney. If your transplanted kidney loses function, dialysis treatment can be started and you can wait for another transplant. (See National Kidney Foundation brochure *Kidney Transplant: A New Lease on Life.*)

What about kidney-pancreas transplants?

Sometimes, if you have Type 1 diabetes, it may be possible to do a pancreas transplant along with a kidney transplant or after it. Your doctor can advise you about this possibility.

What does hemodialysis involve?

Hemodialysis is the most common form of treatment for kidney failure. In order to have hemodialysis, you will need to have surgery to join one of your arteries to a nearby vein in your arm. This will make a bigger vein called a **fistula**. Needles are put into this fistula and joined to the artificial kidney machine. This machine cleans the blood and takes away products that build up in your body. The treatments last about four hours and usually need to be done three times a week. This dialysis can be done in a hospital, an outpatient dialysis clinic or at home (after training). (See National Kidney Foundation brochure, *What You Need to Know About Hemodialysis*.)

What does peritoneal dialysis involve?

Peritoneal dialysis is used often in patients with diabetes. In this type of dialysis, the patient's blood is not cleaned outside the body as with hemodialysis. The blood stays in the blood vessels that line your own abdominal (peritoneal) space. The lining of this space acts like a natural filter.

A plastic tube called a **catheter** is placed into your abdomen by a surgical procedure. During the treatment your abdomen (through the catheter) is slowly filled with fluid called dialysate. Wastes filter out of the blood vessels lining your abdominal space and go into the dialysate fluid. Once the process is finished, the used dialysate is drained out and discarded. This process is then repeated (usually four to six times during the day or night). Several kinds of peritoneal dialysis can be done. (See National Kidney Foundation brochure *What You Need to Know About Peritoneal Dialysis*.)



Key points to remember about diabetes and the kidneys

- ❖ Severe kidney failure may occur in about a third of people with diabetes.
- ❖ Because diabetes may injure the blood vessels in the body, it can cause permanent kidney damage. The earliest sign of kidney damage is an increase in albumin in the urine.
- ❖ The presence of kidney damage and the level of kidney function (GFR) indicate chronic kidney disease.
- ❖ Early kidney damage by diabetes can be found by a sensitive test for a tiny amount of

albumin in the urine (microalbuminuria). A blood test for creatinine may be used to estimate your GFR, which tells how much kidney function you have.

- ❖ Reaching your target blood pressure with the help of ACE inhibitors or angiotensin II receptor blockers is the most effective way of slowing the loss of kidney function.
- ❖ Reducing the amount of salt in your diet may be needed if there is kidney damage, body swelling or high blood pressure.
- ❖ Other factors can cause kidney damage and affect kidney function. These are: blocking of urine flow, urinary tract infections and certain medicines (especially anti-inflammatory pain-relieving medicines).
- ❖ Ankle and leg swelling (edema), going to the bathroom more often at night, high blood pressure and a drop in the amount of insulin needed to control diabetes may all be signs of kidney damage.
- ❖ If chronic kidney disease gets worse and causes kidney failure, this can be treated by kidney transplantation, hemodialysis or peritoneal dialysis. The type of treatment is chosen for every patient, depending on overall health, lifestyle and personal preference. A treatment plan may use each of these forms of therapy at one time or another.
- ❖ The diabetic diet is a very important part of the treatment of all patients with diabetes even if they don't have chronic kidney disease.

Other Resources

You may also be interested in these other publications from the National Kidney Foundation:

- ❖ *Your Kidneys: Master Chemists of the Body*
- ❖ *High Blood Pressure and Your Kidneys*
- ❖ *About Kidney Disease: A Guide for Patients and their Families*
- ❖ *Are You at Increased Risk for Chronic Kidney Disease?*
- ❖ *What You Need to Know About Urinalysis*
- ❖ *Nutrition and Chronic Kidney Disease*
- ❖ *What You Need to Know When You Have Chronic Kidney Disease*
- ❖ *Choosing a Treatment for Kidney Failure*
- ❖ *What You Need to Know About Hemodialysis*
- ❖ *What You Need to Know About Peritoneal Dialysis*
- ❖ *Kidney Transplant: A New Lease on Life.*

To obtain copies of these publications or information about other National Kidney Foundation resources, contact your local National Kidney Foundation affiliate, or call the national toll-free number (800) 622-9010. Also, visit NKF's Web site at www.kidney.org

NOTES

More than 20 million Americans—one in nine adults—have chronic kidney disease, and most don't even know it. More than 20 million others are at increased risk. The National Kidney Foundation, a major voluntary health organization, seeks to prevent kidney and urinary tract diseases, improve the health and well-being of individuals and families affected by these diseases, and increase the availability of all organs for transplantation. Through its 51 affiliates nationwide, the foundation conducts programs in research, professional education, patient and community services, public education and organ donation. The work of the National Kidney Foundation is funded by public donations.

© 1989, 1996, National Kidney Foundation
2002 Edition

All Rights Reserved.

No part of this publication may be reproduced or transmitted in any form or by any means, electronic or mechanical, including photocopy, recording, or any information storage and retrieval system, without permission in writing from the National Kidney Foundation.

- This brochure is also available in Spanish.
- Este folleto se puede conseguir en español.



**National Kidney
Foundation®**

Making Lives Better

National Kidney Foundation
30 East 33rd Street
New York, NY 10016
1-800-622-9010
www.kidney.org

02-09

K/DOQI Learning System (KLS)™