


**National Kidney
Foundation®**



Readers of the **Pocket Guide to Nutrition Assessment of the Patient with Chronic Kidney Disease (4th Edition)** are advised of the following corrections:

Page 1-51

MNT introductory paragraph should read as follows:

CMS covers MNT for two diseases/conditions: 1) Diabetes mellitus Type 1 and 2, or gestational diabetes; and 2) chronic renal insufficiency (non-dialysis) when GFR < 50 mL/min/1.73m² and post-transplant care after discharge from the hospital.

Page 2-8

Significance of Abnormal for carbon dioxide should read: High: Metabolic alkalosis, Low: Metabolic acidosis

Page 4-12

Simple estimate based on dextrose concentration and estimated absorption of 60-76% with CAPD. This should read 60-70%, not 60-76%.

Page 6-13

Enteral Nutrition Supplements: As noted on this list, nutrient content may change regularly and various comparison sites also provide different values for the same products. Manufacturers' Web sites recommend using the product label for the most up-to-date information.

Boost GluControl is now *Boost Glucose Control* and the composition is slightly different than published.

See <http://www.nestle-nutrition.com/Products/Default.aspx>

Nepro Carb Steady: Ca⁺⁺ should read 250, not 260 mg.

See <http://abbottnutrition.com/products/products.aspx?pid=264#factlabel>

Suplena Carb Steady: Protein is 10.6 g, Na⁺ 185 mg, K⁺ 265 mg; other nutrients are as reported when compared to the product web site. See <http://abbottnutrition.com/products/products.aspx?pid=266>

Page 11-23

Protein recommendation for *Immediately Post-Transplant*: 1.3-2 gm/kg/d, not 1.2-3 kg. Protein recommendation for *Long-Term Transplant*: 0.8-1.0 g/kg/d, not 0.8-1.0 g/kg.