

## **TRANSCRIPTIONAL ANALYSIS OF IgA NEPHROPAHY BIOPSIES: TOWARDS A MOLECULAR CLASSIFICATION**

Celine C Berthier<sup>1</sup>, Vidooshi G Maru<sup>1</sup>, Viji Nair<sup>1</sup>, Felix Eichinger<sup>1</sup>, Anna Henger<sup>1</sup>, Anissa Boucherot<sup>1,2</sup>, Maria P. Rastaldi<sup>3</sup>, Hermann-Josef Groene<sup>4</sup>, Clemens D. Cohen<sup>2</sup>, Matthias Kretzler<sup>1</sup> and the European Renal cDNA Bank (ERCB) consortium

<sup>1</sup>Internal Medicine, University of Michigan-Ann Arbor-MI; <sup>2</sup>University of Zurich-Switzerland; <sup>3</sup>Immunopathology-Milan-Italy; <sup>4</sup>German Cancer Research Center-Heidelberg-Germany.

IgA nephropathy (IgAN) is defined by histopathological features. It is currently unclear if a single or multiple pathogenetic mechanisms are activated in IgAN. Genome wide expression profiles of micro-dissected IgAN and living donor (LD) renal biopsies were generated to define IgAN transcriptional-associated pathways and molecular subgroups.

Renal biopsies from LD (n=6) and patients with IgAN (n=27) were collected from an international multi-center cohort. Affymetrix genechips were processed separately from glomeruli and tubulointerstitium, and analyzed using a customized data processing pipeline ([www.GenePattern.org](http://www.GenePattern.org)).

Unsupervised hierarchical clustering segregated controls and IgAN biopsies. Using a Ridge regression analysis, candidate markers defined with a FDR below 5% significantly predicted GFR-MDRD in the tubulointerstitium compartment. Within the IgAN cohort, two subgroups of IgAN patients were defined: one with a fingerprint related to the control group and one with a distinct signature. Analyzing the mRNA and associated pathways responsible for this segregation, Integrin signaling was a top scoring canonical pathway mainly activated in the IgAN group 2 vs. the group 1. GFR-MDRD was significantly lower in group 2 vs. group 1; whereas proteinuria was increased. The segregation observed in IgAN patients was validated by testing a second IgAN cohort (n=14) using a custom array platform.

Gene expression profiles of IgAN biopsies can segregate subgroups based on molecular pathways associated with disease subtypes, which might provide a novel type of information for defining molecular targets and predicting disease course.