

Nutrition and Peritoneal Dialysis









NATIONAL KIDNEY
FOUNDATION®

Stages of chronic kidney disease

There are 5 stages of kidney disease as shown in the table below. Your healthcare provider will tell you the stage of kidney disease, based on how well your kidneys are working and your estimated glomerular filtration rate (eGFR). The eGFR number comes from a lab test that measures the amount of blood your kidneys are filtering each minute. As CKD gets worse, the eGFR number goes down.

STAGES OF KIDNEY DISEASE

STAGE	DESCRIPTION	ESTIMATED GLOMERULAR FILTRATION RATE (eGFR)	KIDNEY FUNCTION
1	Kidney damage (e.g., protein in the urine) with normal kidney function	90 or above	
2	Kidney damage with mild loss of kidney function	60 to 89	
3a	Mild to moderate loss of kidney function	45 to 59	
3b	Moderate to severe loss of kidney function	30 to 44	
4	Severe loss of kidney function	15 to 29	
5	Kidney failure	Less than 15	


Contents

Introduction	4
Getting the right amount of calories	5
Working with your dietitian	6
Getting the right amount of protein.	8
Other important nutrients in your diet.	9
Handling special diet needs	15
How your nutritional health is checked	16
Physical nutrition exam	17
Amount of dialysis you receive.	18
Understanding your test results	19
Where can you get more information?	24
Setting a standard for care.	25



Introduction

If you are receiving peritoneal dialysis treatments, your diet is an important part of your overall care. This booklet will tell you about some things that are important to your diet. It has been written for adults who are receiving peritoneal dialysis treatment. The information is based on recommendations made by the National Kidney Foundation's Kidney Disease Outcomes Quality Initiative. These recommendations have been developed to help your health care team provide the best care for you.



If you are receiving hemodialysis treatment, see *Nutrition and Hemodialysis*. For more information about diets for transplantation, see *Nutrition and Transplantation*. For information about diets for children with chronic kidney disease, see *A Parent's Guide to Nutrition for Children With Chronic Kidney Disease*. All are available from the National Kidney Foundation.

Getting the right amount of calories

Getting the right amount of calories is important to your overall health and how well you feel. Calories come from all the foods you eat. They are important because they:

- Give your body **energy**
- Help you maintain a **healthy weight**
- Help your body use protein for **building muscles and tissues**.

When you first start peritoneal dialysis, you may have difficulty eating well and getting enough calories. For a while, the dialysis solutions may give you a sense of fullness in your stomach. Eating smaller meals five or six times a day can provide the calories you need during the first weeks on peritoneal dialysis.

Over time, many people gain unwanted weight on peritoneal dialysis. The dialysis fluid used for exchanges contains a sugar called dextrose. Solutions that contain more dextrose help to remove extra fluid from your blood.

However, dextrose is an extra source of calories for the body and can lead to unwanted weight gain. And if you have diabetes, the extra sugar from your dialysis solution can cause an increase in your blood sugar. The registered dietitian nutritionist (RDN) at your dialysis center can help you plan meals to prevent extra weight gain and high blood sugar. In addition, following the sodium and fluid instructions from your dietitian can help to prevent the need for the high sugar solutions. Your healthcare professional will choose the dialysis solutions for your fluid removal. Also, your healthcare professional may change your diabetic medicines to help control blood sugar.

Working with your dietitian

You may feel a bit confused by all the new information about your kidney disease and its treatment. You probably have many questions about your diet. Help is available to you. The staff at your dialysis center includes a registered dietitian nutritionist (RDN) with special training in diets for people with kidney disease. This dietitian can answer your questions about your diet and help you plan your meals to get the right foods in the right amounts.



Steps to take

- Speak to the dietitian at your dialysis center.
- Ask your dietitian to **help you plan meals** with the right amount of calories.
- Keep a **diary** of what you eat. Show this to your dietitian on a regular basis.
- Ask your healthcare professional and dietitian what is your **best weight**. Weigh yourself each day in the morning.
- If you are **losing too much weight**, ask your dietitian how to add extra calories to your diet.
- If you are **slowly gaining too much body weight**, ask for suggestions on safely reducing your daily calorie intake and increasing your activity level.
- If you **gain weight rapidly**, speak to your healthcare professional. A sudden increase in weight, along with swelling, shortness of breath and a rise in your blood pressure may be a sign that you have too much fluid in your body.

Getting the right amount of protein

Before you started dialysis, you may have been on a low protein diet to limit the amount of waste products in your blood. Now that you have begun peritoneal dialysis, your treatments will remove these waste products. Unfortunately, when your dialysis removes the unwanted wastes, it also carries out some good proteins that your body needs. Eating a higher protein diet can help you replace the lost protein.

Your body needs the right amount of protein for:

- building muscles
- repairing tissue
- fighting infection

Protein-rich foods to eat daily include:

- fresh meats
- poultry (chicken and turkey)
- fish and seafood
- eggs or egg whites
- small amounts of dairy products

Some of these protein-rich foods may also contain a lot of phosphorus, a mineral you may need to control in your diet. Your dietitian will help you plan the right amount and type of each protein source for good health and strength. (For more information about phosphorus, see page 11.)

Steps to take

- Ask your dietitian how much protein you need to eat each day.
- Show your daily food diary to your dietitian, and ask if you are eating the right amount of protein.

Other important nutrients in your diet

Sodium and fluid

Sodium is a mineral found naturally in foods. It is found in large amounts in table salt and in foods that have added table salt such as:

- salty seasonings like soy sauce, teriyaki sauce and garlic or onion salt
- most canned foods (including canned soups and meats)
- most frozen dinners
- processed meats like ham, bacon, sausage and cold cuts
- salted snack foods like chips and crackers
- canned or dehydrated soups (like ramen noodle soup)
- most restaurant meals.

Eating too much sodium can make you thirsty and cause your body to hold more fluid. The extra sodium and fluid can cause:

- swelling or puffiness around eyes, hands or feet
- fluid weight gain
- shortness of breath
- a rise in blood pressure
- more work for your heart.

Be sure to follow your recommended sodium allowance. Learn to flavor your foods with herbs and spices instead of table salt. **Do not use salt substitutes containing potassium unless approved by your healthcare professional.**



TIP

Try using fresh or dried herbs and spices instead of salt to enhance the flavor of your foods. Also, try adding a dash of hot pepper sauce or a squeeze of lemon juice for flavor.



Phosphorus and calcium

Phosphorus is a mineral found in all foods. Large amounts of phosphorus are found in:

- milk, yogurt and ice cream
- cheese
- nuts and peanut butter
- dried beans and peas
- dark cola drinks
- convenience and processed foods with phosphorus-containing food additives

Eating foods high in phosphorus will raise the amount of phosphorus in your blood. However, phosphorus from plant foods is less absorbed than phosphorus from animal foods or phosphate additives. Dialysis cannot remove all this phosphorus. When phosphorus builds up in the blood, calcium is pulled from the bones. Over time, this may cause bones to become

weak and break easily. It may also cause calcium-phosphorus crystals to build up in your joints, muscles, skin, blood vessels and heart. These deposits may cause serious conditions such as bone pain, organ or heart damage, poor blood circulation and skin infection.

To keep blood phosphorus at safe levels, you will need to **limit phosphorus-rich foods**, and you will need to take a medicine called a **phosphate binder**. These binders are taken with every meal and snack.



TIP

Using fresh foods instead of processed or convenience foods is a good way to lower the amount of phosphorus in your diet.

Foods that are good sources of calcium are also high in phosphorus. The best way to prevent loss of calcium from your bones is to follow a diet that limits high-phosphorus foods and to take phosphate binders. Your healthcare professional may also recommend that you take a special form of vitamin D to help keep calcium and phosphorus in balance and prevent bone disease. Do not take over-the-counter vitamin D, however, unless recommended by your kidney healthcare professional.

Potassium


Potassium is another important mineral found in food. Potassium helps your muscles, and heart, to work properly. Too much or too little potassium in the blood can be dangerous. With peritoneal dialysis, you may need to increase or decrease the amount of potassium in your diet. Each person is different. Your blood level of potassium will be checked every month and your dietitian will help you plan a diet that will give you the right amount of potassium from your foods. If your potassium levels are very low, your healthcare professional may ask you to take a potassium supplement to keep the right amount of potassium in your blood. Large amounts of potassium are found in:

- certain fruits and vegetables (like bananas, oranges, potatoes and some juices)
- milk and yogurt
- dried beans and peas
- most salt substitutes
- protein-rich foods like meats, poultry and fish



Vitamins and minerals

Eating a variety of foods gives your body the vitamins and minerals it needs each day. Your healthcare professional may order special vitamin and mineral supplements for two reasons. Dialysis treatment changes your vitamin needs. Also, your special diet may limit some important food groups. Take only those supplements your kidney healthcare professional orders since **certain vitamins and minerals can be harmful if you are on dialysis**. Also check with your healthcare professional before using any herbal remedies, as some of these may also be harmful for people with kidney disease.



Handling special diet needs

Diabetes and your special diet

You may need to make only a few changes in your diabetic diet to fit your needs on peritoneal dialysis. You may need to eat more protein and fewer carbohydrates. Your dietitian will help develop a meal plan especially for you.

Plant-based diets

Plant-based diets may have a positive effect on health. Eating a variety of plant foods and getting enough calories is important. Eating enough calories is an important way to use protein for important jobs like building muscle, healing wounds and fighting infections. Talk with your dietitian about the best sources of plant protein that have the right amounts of potassium and phosphorus to meet your special needs.



How your nutritional health is checked

There are several different ways for your healthcare professional and dietitian to know if you are eating the right amount of calories or protein. The following sections explain these tests. If your results are not as good as they should be, ask how to improve them. For more information, see “Understanding Your Test Results” at the back of this booklet. You may also want to track your important test results by using the Dialysis Lab Log, available by calling the NKF Cares Patient Help Line at **855.NKF.CARES** (855.653.2273).

Dietary interviews and food diaries

Your dietitian will speak to you at times about your diet. The dietitian may also ask you to keep a record of what you eat each day. If you are not getting enough protein, calories and other nutrients, the dietitian will give you ideas about foods that will improve your diet.



Physical nutrition exam

Your dietitian may use a method called Subjective Global Assessment (SGA) to check your body for signs of nutrition problems. This involves asking you some questions about your daily food intake and checking the fat and muscle stores in your body. The dietitian will consider:

- changes in your weight
- changes in your face, arms, shoulders, hands and legs
- your food intake
- your activity and energy levels
- problems that might interfere with eating

Amount of dialysis you receive

About every three to six months, tests will be performed to see if the amount of dialysis you are getting is enough to keep you in overall good health. The tests include a 24-hour urine collection, samples of your dialysis solution and a blood test. This information will measure the amount of dialysis you receive, called Kt/V (pronounced kay tee over vee). A low Kt/V indicates you are not getting enough dialysis. Low amounts of dialysis can keep you from feeling well, sleeping soundly or eating well. It is very important to do all your dialysis exchanges as ordered by your healthcare professional to keep your Kt/V level as high as possible.

Steps to take

- Ask your healthcare professional and dietitian what tests will be used to check your nutritional health.
- Ask for a copy of the Dialysis Lab Log and track your results.
- If your numbers are not in the normal range, ask your healthcare professional and dietitian how you can improve them.



Understanding your test results

Some or all of the following tests may be used to check your nutrition and general health. Ask your healthcare professional and dialysis care team which tests you will have, how often they will be done and what your levels should be. If your numbers are not in range, ask how to improve them.

Blood Pressure: If your blood pressure is high, make sure to follow all the steps in your prescribed treatment. These steps may include taking high blood pressure medicines, cutting down on the amount of salt in your diet, losing weight if you are overweight and following a regular exercise program.

Blood UreaNitrogen (BUN): Urea nitrogen is a normal waste product in your blood that comes from the breakdown of protein from foods you eat. Healthy kidneys remove BUN from your blood, but when kidney failure occurs, your BUN rises. BUN is also removed from your blood by your dialysis. Your BUN rises from not getting enough dialysis or from

eating too much protein. It can fall from getting more dialysis or from eating the right amount of protein recommended by your healthcare professional and dietitian.

Body Weight: Maintaining a healthy weight is important to your overall health. If you are losing weight without even trying, you may not be getting the right nutrition to stay healthy. Your dietitian can suggest how to safely add extra calories to your diet. On the other hand, if you are slowly gaining unwanted weight, you may need to eat less calories and increase your activity level. A sudden weight gain may also be a problem. If it is accompanied by swelling, shortness of breath and a rise in blood pressure, it may be a sign of too much fluid in your body. You should check your weight at home every morning. Speak to your healthcare professional if your weight changes suddenly.

Calcium: Calcium is a mineral that is important for strong bones. To help balance the amount of calcium in your blood, you should avoid foods fortified with calcium and do not take a calcium supplement unless recommended by your healthcare professional. Your healthcare professional may also prescribe a special form of vitamin D. Take only the medicines recommended by your healthcare professional.

Cholesterol:

- **Total:** Cholesterol is a fat-like substance found in your blood. A high cholesterol level may increase your risk of having heart and circulation problems. However, a cholesterol level that is too low may mean you are not eating well enough to stay healthy.
- **HDL:** HDL cholesterol is a type of “good” cholesterol that protects your heart.
- **LDL:** LDL cholesterol is a type of “bad” cholesterol. A high LDL level may increase your chance of having heart and circulation problems. If your LDL level is too high, your healthcare professional may recommend changing your diet and increasing your activity level.

Serum Creatinine: Creatinine is a waste product in your blood that comes from the normal function of your muscles. Healthy kidneys remove creatinine from your blood, but when the kidneys are not working, your creatinine level rises. Your dialysis also removes creatinine from your blood. Not getting enough dialysis can cause your creatinine level to rise, while getting more dialysis causes it to fall. Your creatinine level can also fall from not eating well over a long period of time.

Hemoglobin: Hemoglobin is the part of red blood cells that carries oxygen from your lungs to all the tissues in your body. Measuring your hemoglobin level tells your healthcare professional if you have anemia, which makes you feel tired and have little energy. To treat your anemia, you may need to take a hormone called ESA along with iron.

Iron:

- **TSAT and Serum Ferritin:** Your TSAT (pronounced tee sat) and serum ferritin (pronounced ferry tin) are measures of iron in your body. They help you build red blood cells. Your healthcare professional will recommend iron when needed to reach your target levels.

Kt/V: Kt/V (pronounced kay tee over vee) is a measure of the amount of dialysis you receive. Getting the right amount of dialysis is important to your overall health and can also affect how well you eat.

Parathyroid Hormone (PTH): High levels of parathyroid hormone (PTH) may result from a poor balance of calcium and phosphorus in your blood. This can cause bone disease. Your healthcare professional may order a special form of vitamin D to help lower your PTH. Caution: Do not take over-the-counter vitamin D unless ordered by your kidney healthcare professional.

Phosphorus: A high phosphorus level in your blood can lead to weak bones, itching, bone pain and hardening of blood vessels. If your level is too high, your healthcare professional may ask you to reduce your intake of foods that are high in phosphorus and take a phosphate binder with all your meals and snacks.

Potassium: Potassium is a mineral that helps your heart and muscles work properly. A potassium level that is too high or too low may weaken muscles and change your heartbeat. Whether you need to change your intake of high-potassium foods varies with each person on peritoneal dialysis. Your dietitian can help you plan your meals to get the right amount of potassium.

Subjective Global Assessment (SGA): Your dietitian may use SGA to help check for signs of nutrition problems. The dietitian will ask you some questions about your daily diet and check your weight and the fat and muscle stores in your face, hands, arms, shoulders and legs. If your SGA score is too low, ask how to improve it.

Triglyceride: Triglyceride is another type of fat found in your blood. A high triglyceride level, along with high levels of total and LDL cholesterol, may increase your chance of having heart and circulation problems.

Where can you get more information?

If you have questions, speak with your healthcare team. They know you and can answer questions about you.

If you want to read more about kidney disease, the National Kidney Foundation has more than 50 other publications that cover many subjects, such as:

- *CKD risk factors like hypertension and diabetes*
- *Complications of chronic kidney disease such as cardiovascular disease, anemia, or bone problems*
- *Nutrition for CKD patients with information about carbohydrates, protein, sodium, phosphorus, and potassium*
- *Treating kidney disease early*
- *Treating kidney failure with transplantation or dialysis transplantation or dialysis.*

There are two ways to learn about the many free resources available to you:

- Call the NKF Cares Patient Help Line toll free at **855.NKF.CARES** (855.653.2273).
- Visit the National Kidney Foundation website at **[kidney.org/store](https://www.kidney.org/store)**.

Becoming an educated patient is very important to being healthy!

Setting a standard for care

The National Kidney Foundation, through its *Kidney Disease Outcomes Quality Initiative* (KDOQI®), defines stages of kidney disease and offers guidelines that help your healthcare professional and healthcare team make important decisions about your medical treatment.

The information in this booklet is based on those recommended guidelines.



The information contained in this publication is based on current data and expert guidance available at the time of publication. The information is intended to help patients become aware of their disease and its management. This publication is not intended to set out a preferred standard of care and should not be construed as one. Neither should the information be interpreted as prescribing an exclusive course of management. Patients should always consult with their healthcare professional regarding decisions about their individual plan of care.



NATIONAL KIDNEY FOUNDATION®

Fueled by passion and urgency, the National Kidney Foundation (NKF) is a lifeline for all people affected by kidney disease. As pioneers of scientific research and innovation, we focus on the whole patient through the lens of kidney health. Relentless in our work, we enhance lives through action, education, and accelerating change.

30 E. 33RD ST. | NEW YORK, NY 10016 | 800.622.9010 | [KIDNEY.ORG](https://www.kidney.org)

KIDNEY.ORG

