

# HIGH BLOOD PRESSURE AND CHRONIC KIDNEY DISEASE

*For People with CKD Stages 1-4*



National  
Kidney  
Foundation®

[www.kidney.org](http://www.kidney.org)

# National Kidney Foundation's Kidney Disease Outcomes Quality Initiative

Did you know that the National Kidney Foundation's Kidney Disease Outcomes Quality Initiative (NKF-KDOQI™) has guidelines and commentaries that help your doctor and healthcare team make important decisions about your medical treatment? The information in this booklet is based on the NKF-KDOQI recommended guidelines and commentaries.

## What is your stage of kidney disease?

There are five stages of kidney disease. They are shown in the table below. Your doctor determines your stage of kidney disease based on the presence of kidney damage and your glomerular filtration rate (GFR), which is a measure of your level of kidney function. Your treatment is based on your stage of kidney disease. Speak to your doctor if you have any questions about your stage of kidney disease or your treatment.

### STAGES OF KIDNEY DISEASE

| Stage | Description  | Glomerular Filtration Rate (GFR)* |
|-------|--|-----------------------------------|
| 1     | Kidney damage (e.g., protein in the urine) with normal GFR | 90 or above                       |
| 2     | Kidney damage with mild decrease in GFR                    | 60 to 89                          |
| 3     | Moderate decrease in GFR                                   | 30 to 59                          |
| 4     | Severe reduction in GFR                                    | 15 to 29                          |
| 5     | Kidney failure   | Less than 15                      |

\*Your GFR number tells your doctor how much kidney function you have. As chronic kidney disease progresses, your GFR number decreases.

## TABLE OF CONTENTS

|  |    |
|--|----|
| Did you know? . . . . .  | 4  |
| What is chronic kidney disease? . . . . .  | 5  |
| What is high blood pressure? . . . . .   | 6  |
| How are high blood pressure and kidney<br>disease related? . . . . .                     | 6  |
| How do I know if my blood pressure<br>is too high? . . . . .                             | 7  |
| How is blood pressure measured?<br>How often should it be checked? . . . . .             | 8  |
| I have high blood pressure but am not<br>sure if I have CKD. What should I do? . . . . . | 9  |
| What tests will I need if I have CKD? . . . . .  | 9  |
| How often will I need to visit my<br>doctor or clinic? . . . . .                         | 10 |
| What will my treatment for high blood pressure<br>and CKD include? . . . . .             | 11 |
| What kinds of changes will<br>I need to make to my diet? . . . . .                       | 11 |
| Will I need to make other changes<br>in my lifestyle? . . . . .                          | 13 |
| What about medications? . . . . .  | 14 |
| What if I have CKD and diabetes? . . . . .   | 15 |
| What if I have a kidney transplant? . . . . .  | 15 |
| What can I do to help control my<br>blood pressure? . . . . .                            | 16 |
| What can my family do? . . . . .   | 17 |
| Key Points to Remember . . . . .   | 18 |
| Where can I get more information? . . . . .  | 19 |
| Test Your Knowledge . . . . .  | 20 |



### Did you know?

- More than half of people with chronic kidney disease (CKD) have high blood pressure.
- High blood pressure increases the chance that kidney disease will get worse.
- High blood pressure makes you more likely to develop heart disease.
- Following your treatment plan carefully and keeping your blood pressure under control can help prevent these complications.
- Treatment for high blood pressure and CKD includes following a healthy diet, exercising, and taking medications.

- If you have kidney disease from diabetes and if you have protein in your urine, the best blood pressure medications for your treatment are angiotensin converting enzyme (ACE) inhibitors or angiotensin receptor blockers (ARBs).

This booklet will answer your questions about high blood pressure and kidney disease and tell you how to stay as healthy as possible when you have these conditions. This booklet is written for people who have high blood pressure and CKD in stages 1-4. If you do not know your stage of kidney disease, see the chart on the inside front cover of this booklet and speak to your doctor.

### **What is chronic kidney disease?**

Chronic kidney disease means the kidneys have been damaged by conditions such as diabetes, high blood pressure, glomerulonephritis (a group of diseases that cause damage to the kidney's filtering units), or other conditions (heredity, kidney stones, etc.). As a result, the kidneys are less able to do the following jobs to help maintain health:

- Remove wastes and extra fluid from your body
- Release hormones that help to:
  - Control blood pressure
  - Maintain strong bones
- Prevent anemia by increasing the number of red blood cells in your body
- Keep the right balance of important chemicals in your blood, such as sodium, potassium, phosphorus, and calcium.

When your kidneys are not working well, wastes can build to high levels in your blood and make you feel sick. Even before that, you may develop complications like high blood pressure, anemia (low red blood cell count), weak bones, poor nutritional health, and nerve damage. Kidney disease also makes you more likely to develop heart and blood vessel disease. These problems may happen slowly, over a long period of time, often without symptoms. CKD may eventually lead to kidney failure, and needing dialysis or a kidney transplant to stay alive. Early detection and treatment can prevent or delay these conditions.

### **What is high blood pressure?**

Blood pressure is the force of your blood against the walls of your blood vessels as your heart pumps blood around your body. If this pressure becomes too high, you are said to have high blood pressure or hypertension.

### **How are high blood pressure and kidney disease related?**

They are related in two ways:

1. High blood pressure is a leading cause of CKD. Over time, high blood pressure can damage blood vessels throughout your body. This can reduce the blood supply to important organs like the kidneys. High blood pressure also damages the tiny filtering units in your kidneys. As a result, the kidneys may stop removing wastes and extra fluid from your blood.

The extra fluid in your blood vessels may build up and raise blood pressure even more.

2. High blood pressure can also come from CKD itself. Your kidneys help control your blood pressure. Diseased kidneys are less able to help regulate blood pressure. As a result, blood pressure increases.

If you have CKD, high blood pressure makes it more likely that your kidney disease will get worse and you will have heart problems. Following your treatment plan and keeping your blood pressure controlled can help keep your kidney disease from getting worse and prevent heart disease.

### **How do I know if my blood pressure is too high?**

The only way to tell if your blood pressure is too high is to have it measured. High blood pressure usually causes no symptoms. That is why it has been called a “silent killer.” A single high reading may not mean you have high blood pressure. It should be confirmed on followup visits to your doctor or clinic.

Blood pressure is measured as two numbers. The top number, or "systolic" blood pressure, is the pressure when your heart is beating. The bottom number, or "diastolic" blood pressure, is the pressure when your heart is resting between beats. A blood pressure reading of 140/90 is read as 140 over 90.

Normal blood pressure in adults 18 and older is less than 120/80. People who have a top number between 120 and 139, or a bottom number between 80 and 89, may be more likely to develop high blood pressure. They will need to take steps to prevent it. In general, blood pressure that stays at 140/90 or higher is considered high.





## **How is blood pressure measured? How often should it be checked?**

Blood pressure is usually checked by using a blood pressure cuff around your arm. It should be checked every time you visit your doctor or clinic. You can also be taught to check your own blood pressure at home. Keep a daily record of your blood pressure and show it to your doctor each visit.



## I have high blood pressure but am not sure if I have CKD. What should I do?

Everyone who has high blood pressure has a greater chance of developing CKD. You should ask your doctor about getting these tests:

- A blood test for creatinine, a waste product from muscle breakdown. This should be used to calculate your "glomerular filtration rate," or GFR. Your GFR measures your level of kidney function. If your GFR is too low, it may mean your kidneys are not able to remove enough wastes and extra fluid from your blood.
- A urine test for protein. Protein in the urine is a sign of kidney damage. A higher urine protein means there is a greater chance your kidney disease may get worse and you may develop heart disease.

## What tests will I need if I have CKD?

In addition to checking your GFR and urine protein, you should get the following:

- Diagnosis of the type of CKD you have. This may include an ultrasound to get a picture of your kidneys to check for any problems.
- Some tests for heart and blood vessel disease. These tests will most likely include:
  - An electrocardiogram (EKG)
  - A blood test for glucose (sugar)
  - A blood test for lipids (fatty substances in your blood, such as cholesterol)
  - Height and weight measurement to find your body mass index (BMI). This tells if you are overweight.

Your doctor will also talk to you about:

- Any side effects from your medications
- Any problems in following your treatment plan

### **How often will I need to visit my doctor or clinic?**

Once you are stable with your treatment, you may not need to visit your doctor or clinic as often. Your doctor will want to see you more often if:

- You start a new medication
- Your medication dose is changed
- Your kidney disease is getting worse
- Your blood pressure rises

At followup visits, your doctor will check:

- Your blood pressure
- Your GFR
- Your urine protein level
- Your blood level of potassium. If your kidney function gets worse, potassium may build up in your blood. This can be dangerous for your heart. Some of the best medicines for treating high blood pressure and preventing further kidney damage can also make the potassium go higher. If your potassium level gets too high, you will need to make some changes in your diet (see "What kinds of changes will I need to make to my diet?" on pages 11–13).

## What will my treatment for high blood pressure and CKD include?

A treatment plan should be developed for you based on your stage of CKD. Your doctor may recommend that you see another doctor with special training in kidney disease or high blood pressure. This doctor will help with your treatment plan. The goals of your treatment are:

- To control your blood pressure
- To keep kidney disease from getting worse
- To lower your chance of getting heart disease

To help reach these goals, you will need to make some lifestyle changes, such as following a healthier diet, exercising more, and taking medications.

## What kinds of changes will I need to make to my diet?

That will depend on your stage of CKD. In stages 1–2, your doctor may want you to follow a diet, such as DASH (Diet Approaches to Stop Hypertension), that is high in fruits, vegetables, and dairy foods. DASH is low in sugar and sodium. If you have any questions about your diet, speak to your doctor. Your doctor can refer you to a registered dietitian who will help you work out a diet plan to meet your needs.



People with CKD in stages 1–4 may also be asked to:

- Cut back on sodium (found in large amounts in table salt and foods with added salt).
- Cut down on foods that are high in saturated fats and cholesterol. These can clog up your arteries and increase your risk for heart and blood vessel disease.
- Control the amount of carbohydrates in your diet. Carbohydrates are foods that turn to sugar when you digest them. They are found in many foods, such as bread, rolls, tortillas, rice, pasta, potatoes, corn, dried beans, fruits, fruit juices, milk, and yogurt.

If you have CKD in stages 3–4, you may be asked to make some other diet changes to keep CKD from getting worse and to prevent complications such as bone disease. You may need to:

- Control the amount of protein you eat. Ask your doctor how much protein you need each day. Your dietitian can help you plan your meals to get the right amount of protein.
- Lower the potassium in your diet. Your doctor will check the amount of potassium in your blood. Ask your doctor if your potassium level is too high. If it is, you will need to lower your intake of potassium-rich foods. Your dietitian can teach you which foods are high and low in potassium, and help you plan your meals to get the right amount.
- Eat fewer foods that are high in phosphorus. High levels of phosphorus in your blood may lead to bone disease. Phosphorus is found in large amounts in dairy products like milk, cheese, pudding, yogurt, and ice cream; in nuts and peanut butter; and in drinks like cocoa, beer, and dark cola.

### **Will I need to make other changes in my lifestyle?**

Yes. The following steps can also help you reach your treatment goals:

- Lose weight if you are overweight.
- Exercise for 30 minutes a day on most days of the week. Always check with your doctor before starting an exercise program.
- Cut back on alcohol (no more than two drinks a day for men, and one drink a day for women).
- Stop smoking if you are a smoker.



## What about medications?

To reach your target blood pressure, you will probably need more than one type of blood pressure medication. Your medications may include:

- An angiotensin-converting enzyme (ACE) inhibitor or an angiotensin receptor blocker (ARB). Studies have shown that these may help protect your kidney function and lower your risk for heart disease if you have protein in your urine.
- A diuretic or water pill to help your body get rid of extra fluid.
- Other medications, such as beta blockers or calcium channel blockers, to help reach your treatment goals.

Be sure to take your medications exactly as prescribed by your doctor. Report any side effects to your doctor right away. It may be possible to lower your dose or switch you to another medication. Do not stop taking any of your medications on your own without letting your doctor know. Follow up to check your blood pressure regularly. Even though you are feeling well, uncontrolled high blood pressure could still damage vital organs like the heart and kidneys. Remember, high blood pressure is a "silent killer."

### **What if I have CKD and diabetes?**

Controlling your blood sugar and blood pressure are the most important things you can do to keep kidney disease from getting worse and to prevent other problems. Be sure to make the lifestyle changes your doctor recommends and take your medications *exactly* as prescribed. If you have diabetes and CKD, an ACE inhibitor or ARB may be part of your treatment plan. This is because these medications can help to protect your kidneys.

### **What if I have a kidney transplant?**

Many kidney transplant recipients have high blood pressure and CKD. It's very important to lower your blood pressure to keep your kidney transplant working well. If you have high blood pressure, be sure to follow the lifestyle changes your doctor recommends and to take your blood pressure medications *exactly* as prescribed.



## What can I do to help control my blood pressure?

You can do a lot.

- Be sure to keep all your appointments with your doctor.
- Ask your doctor how you can learn to take your own blood pressure at home. Keep a daily record of your blood pressure and show it to your doctor each visit.
- Take your blood pressure pills *exactly* as prescribed by your doctor, even if you are feeling fine. High blood pressure usually causes no symptoms.
- Report any side effects from your medicine to your doctor. Never stop taking any of your medications on your own.
- To help you remember when to take your medications, try one of the special pill boxes with small compartments labeled with the days of the week and the time of day. Cellphones or watches with alarms can also help. Ask your family members to remind you to take your medications.
- Follow your diet and exercise plan faithfully. If you need help with your diet, ask your doctor to refer you to a registered dietitian. The dietitian will explain the diet and help you develop a meal plan you can live with.
- Learn all you can about your treatment. Don't be afraid to ask questions if there is something you don't understand. Bring a list of your questions when you go to the doctor.

## What can my family do?

It's a good idea to get your whole family involved in your care. It can be much easier to make lifestyle changes, such as following a healthier diet, exercising more, and stopping smoking, if you have the support of your family. Because high blood pressure often runs in families, some of your family members may also be at risk for high blood pressure and chronic kidney disease (CKD). You should tell them to learn all they can about high blood pressure and to have their blood pressure checked at least once a year.



## Key Points to Remember

- High blood pressure and kidney disease are closely related. High blood pressure is a cause of kidney disease. It can also come from having kidney disease.
- Having high blood pressure increases the chance that kidney disease will get worse and that heart problems will develop.
- Keeping blood pressure well controlled reduces the chances of these health problems.
- High blood pressure usually causes no symptoms. The only way to find out if your blood pressure is too high is to have it measured. Your blood pressure should be checked at every visit to your doctor or clinic.
- If you have CKD, your doctor should also test your blood to check your kidney function (GFR) and for levels of other chemicals in your blood such as potassium. Your urine will be checked by your doctor for protein, a sign of kidney problems.
- Your treatment will include making changes for a healthier lifestyle and taking medications.
- Your treatment goals are: control your blood pressure to keep kidney disease from getting worse and reduce your chance of getting heart disease.
- You will probably need more than one type of high blood pressure pill to reach your target. These may include: a diuretic or water pill, an ACE inhibitor or ARB, and other medications.
- Report any problems to your doctor. Never stop taking a medication without speaking to your doctor.
- Learn all you can about high blood pressure and CKD.

- Get involved in your own care.
- Encourage family members to have their blood pressure checked. They may also be at increased risk for high blood pressure and CKD.

## Where can I get more information?

If you have questions, speak with your healthcare team. They know you and can answer questions about you.

If you want to read more about kidney disease, the National Kidney Foundation has more than 50 publications that cover many subjects, such as:

- CKD risk factors like high blood pressure and diabetes
- Complications of CKD, such as cardiovascular disease, anemia, or bone problems
- Nutrition for CKD patients; information about carbohydrates, protein, sodium, phosphorus, and potassium
- Treating kidney disease early
- Treating kidney failure with transplantation or dialysis

There are two ways to learn about the many free resources available to you:

- Call the National Kidney Foundation at 855.NKF.CARES (855.653.2273) or email **NKFCares@kidney.org**
- Visit the National Kidney Foundation website: **www.kidney.org/store**. (All publications are free, but there is a limit of five per person.)

Becoming an educated patient is very important to being healthy!

## Test Your Knowledge: Take This True or False Quiz.

1. High blood pressure increases the chance that kidney disease will get worse.  
True \_\_\_\_\_ False \_\_\_\_\_
2. High blood pressure may be a complication of kidney disease.  
True \_\_\_\_\_ False \_\_\_\_\_
3. High blood pressure causes many symptoms.  
True \_\_\_\_\_ False \_\_\_\_\_
4. The treatment for high blood pressure includes lifestyle changes and medications.  
True \_\_\_\_\_ False \_\_\_\_\_
5. Usually, one medication is enough to control high blood pressure in people with CKD.  
True \_\_\_\_\_ False \_\_\_\_\_
6. Getting regular exercise may help reduce your blood pressure.  
True \_\_\_\_\_ False \_\_\_\_\_
7. If you have high blood pressure, you may need to increase sodium in your diet.  
True \_\_\_\_\_ False \_\_\_\_\_
8. If you have kidney disease, you are also at higher risk for heart disease.  
True \_\_\_\_\_ False \_\_\_\_\_
9. High blood pressure often runs in families.  
True \_\_\_\_\_ False \_\_\_\_\_
10. You can't check your own blood pressure.  
True \_\_\_\_\_ False \_\_\_\_\_

See answers on page 21.

## Questions for My Doctor

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

### Answers to Quiz on page 20

- |      |       |
|------|-------|
| 1. T | 6. T  |
| 2. T | 7. F  |
| 3. F | 8. T  |
| 4. T | 9. T  |
| 5. F | 10. F |

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---





**The National Kidney Foundation** is the leading organization in the U.S. dedicated to the awareness, prevention, and treatment of kidney disease for hundreds of thousands of healthcare professionals, millions of patients and their families, and tens of millions of Americans at risk.



National  
Kidney  
Foundation®

30 East 33rd Street  
New York, NY 10016

---

Awareness. Prevention. Treatment.

[www.kidney.org](http://www.kidney.org) 800.622.9010

