


## Past KDOQI Poll Questions and Results

April 2017 Poll

Q1 Should patients with CKD stage 3-5 (non-dialysis) and hyperphosphatemia receive non-calcium containing binders only?


Answered: 974 Skipped: 33



ANSWER CHOICES	RESPONSES	
Yes	46.00%	448
No	54.00%	526
TOTAL		974

Q2 Should patients with hypertension who are on dialysis preferentially receive ACE inhibitors/ARBs (as opposed to beta blockers)?


Answered: 979 Skipped: 28



ANSWER CHOICES	RESPONSES	
Yes	68.23%	668
No	31.77%	311
TOTAL		979

**Q3 Should statin medications be discontinued in patients with CKD 5D who are age 75 years and older?**

Answered: 980 Skipped: 27




ANSWER CHOICES	RESPONSES	
Yes	46.53%	456
No	53.47%	524
<b>TOTAL</b>		<b>980</b>

January 2017 Poll

**Q1 Should Rituximab be considered first-line treatment of membranous nephropathy, assuming a clinical indication for immunosuppressant therapy?**


Answered: 682 Skipped: 58



Answer Choices	Responses	
Yes	49.85%	340
No	50.15%	342
<b>Total</b>		<b>682</b>

**Q2 Should systolic blood pressure be lowered to <120 mmHg in patients age 50 years and older with stage 3-5 non-dialysis dependent kidney disease, in the absence of diabetes and/or overt proteinuria?**


Answered: 687 Skipped: 53



Answer Choices	Responses	
Yes	39.01%	268
No	60.99%	419
<b>Total</b>		<b>687</b>

**Q3 Should ACE inhibitors or angiotensin receptor blockers be discontinued in patients with non-dialysis dependent CKD once eGFR drops below 20?**

Answered: 703 Skipped: 37




Answer Choices	Responses	
Yes	46.23%	325
No	53.77%	378
<b>Total</b>		<b>703</b>

September 2016 Poll

**Q1 Should patients with non-dialysis dependent CKD, proteinuria, and inability to take ACEi or ARB due to hyperkalemia, be placed on chronic K+ binding medications plus ACEi or ARB?**


Answered: 189 Skipped: 6



Answer Choices	Responses	
Yes	53.97%	102
No	46.03%	87
<b>Total</b>		<b>189</b>

**Q2 Should patients with significant residual renal function starting hemodialysis be started on 2x per week therapy?**


Answered: 189 Skipped: 6



Answer Choices	Responses	
Yes	58.20%	110
No	41.80%	79
<b>Total</b>		<b>189</b>

**Q3 Should older patients (age >75 years) with co-morbidities approaching dialysis, who opt for hemodialysis, be referred ahead of time for an arteriovenous fistula or referred “just in time” for arteriovenous graft?**

Answered: 191 Skipped: 4




Answer Choices	Responses
Fistula (with earlier referral)	78.53% 150
Graft	21.47% 41
<b>Total</b>	<b>191</b>

June 2016 Poll

**Q1 Should obese patients receiving dialysis be required to lose weight before being actively wait-listed for a kidney transplant?**


Answered: 3,424 Skipped: 90



Answer Choices	Responses	
Yes	70.82%	2,425
No	29.18%	999
<b>Total</b>		<b>3,424</b>

**Q2 Should warfarin be used in dialysis patients with atrial fibrillation?**


Answered: 3,074 Skipped: 440



Answer Choices	Responses	
Yes	54.85%	1,686
No	45.15%	1,388
<b>Total</b>		<b>3,074</b>

### Q3 Should diuretics be used in dialysis patients with urine output?

Answered: 3,203 Skipped: 311



Answer Choices	Responses	
Yes	59.51%	1,906
No	40.49%	1,297
<b>Total</b>		<b>3,203</b>