



Vaccination Schedule for Children and Teens With Chronic Kidney Disease or Kidney Transplant

Age	HepB Hepatitis B	DTaP/Tdap Diphtheria, tetanus, pertussis (whooping cough)	Hib <i>Haemophilus influenzae</i> type b	IPV Polio	PCV13 Pneumococcal conjugate	RV Rotavirus	MMR Measles, mumps, rubella	Varicella Chickenpox	HepA Hepatitis A	HPV Human papillomavirus	Men- ACWY	MenB	Influenza Flu
											Meningococcal		
Children who have an organ transplant or are immune-suppressed should not receive HBV, MMR, or Varicella													
Birth	✓												
2 months	✓	✓	✓	✓	✓	✓							
4 months	✓	✓	✓	✓	✓	✓							
6 months	✓	✓	✓	✓	✓	✓ ¹							✓
12 months	(6–18 mos)	✓ ¹	✓	(6–18 mos)	✓		✓	✓	✓✓				(One dose each fall or winter to all people ages 6 mos and older. Some children younger than age 9 years need 2 doses; ask your child's healthcare provider if your child needs more than 1 dose.)
15 months		✓	(12–15 mos)		(12–15 mos)								
18 months		(15–18 mos)											
19–23 months													
4–6 years		✓		✓			✓	✓					
7–10 years													
11–12 years		✓ (Tdap)								✓✓✓	✓		
13–15 years													
16–18 years											✓	✓✓✓ ²	

¹ This dose of DTaP may be given as early as age 12 months if it has been 6 months since the previous dose. ² Your child may not need this dose depending on the type of vaccine that your healthcare provider uses.