Are You at Increased Risk for Chronic Kidney Disease?







National Kidney Foundation's Kidney Disease Outcomes Quality Initiative

Did you know that the National Kidney Foundation's Kidney Disease Outcomes Quality Initiative (NKF-KDOQI)™ develops guidelines that help your doctor and healthcare team make important decisions about your medical treatment? The information in this booklet is based on the NKF-KDOQI recommended guidelines.

Stages of Chronic Kidney Disease

There are five stages of chronic kidney disease. They are shown in the table below. When you have kidney disease, your doctor determines your stage of kidney disease based on the presence of kidney damage and your glomerular filtration rate (GFR). GFR is a measure of your level of kidney function. Treatment is based on the stage of kidney disease.

Stages of Kidney Disease					
Stage	Description	Glomerular Filtration			
		Rate (GFR)*			
1	Kidney damage (e.g., protein				
	in the urine) with normal GFR	90 or above			
2	Kidney damage with mild				
	decrease in GFR	60 to 89			
3	Moderate decrease in GFR	30 to 59			
4	Severe reduction in GFR	15 to 29			
5	Kidney failure	Less than 15			
*Your GFR number tells your doctor how much kidney function					

^{*}Your GFR number tells your doctor how much kidney function you have. As chronic kidney disease progresses, your GFR number decreases.

Table of Contents

to good health?
What is chronic kidney disease? 6
How do I know if I am at increased risk for chronic kidney disease? 8
Why are African Americans and other ethnic groups at increased risk for chronic kidney disease? 9
What should I do if I am at increased risk for chronic kidney disease? 9
What are the symptoms of chronic kidney disease?
What happens if my test results show I may have chronic kidney disease? 12
What should I do if I am at increased risk but I do not yet have chronic kidney disease?
If I have chronic kidney disease, can I keep it from getting worse? 15
How is chronic kidney disease treated?. 16
What happens if kidney failure occurs? . 20
Is there anything else I should know? 21
Key points to remember 22
What other information is available? 24

Did you know that...

- Having chronic kidney disease (CKD) increases your chance of premature death. This is because people with CKD are more likely to develop heart and blood vessel disease.
- Cardiovascular disease is the number one cause of death among kidney patients.
- Your doctor should order some simple tests to see if you have CKD. People with diabetes or hypertension are at increased risk. There are also other risk factors for CKD.
- More than 26 million American adults have CKD, and most don't even know it. Millions of others are at increased risk for developing CKD and its complications.
- Early detection and treatment can keep kidney disease from getting worse and prevent complications like kidney failure, heart disease and premature death.

This booklet will help you learn what to do about your increased risk of kidney problems.

Why are the kidneys so important to good health?

Your kidneys do some important jobs to keep your body healthy. They:

- Balance your body fluids. Excess fluid is filtered out of your blood, and it leaves your body as urine. Your kidneys make about one to two quarts of urine a day.
- Regulate body water and important minerals in your blood, such as sodium, potassium, phosphorus and calcium.
- Remove waste products from your blood. These waste products come from the breakdown of foods you eat and from normal muscle activity.
- Remove drugs and toxins from your body.
- Release hormones into your blood, which:
 - control blood pressure
 - make red blood cells
 - keep your bones healthy.

What is chronic kidney disease?

Chronic kidney disease (CKD) is a condition that damages your kidneys. It decreases your kidneys' ability to do the jobs above and keep you healthy. The two main causes of CKD are diabetes and high blood pressure. Diabetes occurs when your blood sugar is too high, causing damage to many organs in your body, including the kidneys and heart as well as blood vessels, eyes and nerves. High blood pressure, or hypertension, occurs when the pressure of blood against the walls of your blood vessels increases. If uncontrolled, or poorly controlled, high blood pressure can be a leading cause of CKD, heart attacks and strokes. CKD can also cause high blood pressure.

Many other conditions can harm the kidneys. These include:

- heart and blood vessel disease
- glomerulonephritis, a disease that causes inflammation in the kidneys

inherited diseases, like polycystic kidney disease, which causes cysts to form in the kidneys.

If CKD gets worse, waste products and fluid may build to high levels in your blood and make you feel sick. You may experience other problems like high blood pressure, anemia, weak bones, poor nutritional health and nerve damage. Anemia means there is a short supply of red blood cells in the body, which can make you feel tired and have little energy.

of worsening heart and blood vessel disease. CKD may progress slowly over a long time. In fact, many people don't even know they have kidney disease until it is severe. If it is found and treated early, CKD may often be slowed or stopped. If it keeps getting worse, however, it may lead to kidney failure. This means your kidneys no longer work well enough to keep you alive, and you need a treatment like dialysis or a kidney transplant.

How do I know if I am at increased risk for chronic kidney disease?

Your doctor or clinic should check to see if you have any risk factors for CKD. These include:

- diabetes
- high blood pressure
- heart or blood vessel problems
- a family history of chronic kidney disease
- older age (65 and older).

CKD is also more common in African Americans, Hispanic Americans, Asian or Pacific Islanders and American Indians.



Why are African Americans and other ethnic groups at increased risk for chronic kidney disease?

Diabetes, the leading cause of CKD, is more common in these groups. Also, high blood pressure, the second leading cause of CKD, appears more often in African Americans than in other ethnic groups. Many experts believe these groups may have an inherited tendency to develop these diseases. When combined with other things such as being overweight, this tendency may lead to disease. Staying at a normal weight and getting enough exercise are very important for these groups to help prevent diabetes and high blood pressure.

What should I do if I am at increased risk for chronic kidney disease?

You should visit your doctor or clinic and get tested. Your checkup should include:

Having your blood pressure checked.

- Having a simple test for protein in your urine. Protein is an important building block in your body. Any filtered protein is normally reabsorbed and kept in your body. When your kidneys are damaged, however, protein leaks into your urine. There are different tests to find protein in your urine. If you have two positive tests over several weeks, you are said to have persistent protein in your urine. This is a sign of CKD.
- Having a simple blood test for creatinine, a waste product that comes from muscle activity. The results of your blood creatinine test should be used to estimate your glomerular filtration rate, or GFR. Your GFR tells how much kidney function you have. A low level of GFR may mean your kidneys are

Tip

The National Kidney
Foundation provides
a free communitybased health program
called the Kidney Early
Evaluation Program
(KEEP ®)

no longer working as well as they should to remove wastes from your body. The National Kidney Foundation (NKF) provides a free community-based health program called the Kidney Early Evaluation Program (KEEP®). This includes tests for the early detection of CKD. Call your local NKF affiliate or the national toll-free number (800.622.9010) to find out if this program is available in your community.

What are the symptoms of chronic kidney disease?

Most people do not have severe symptoms until the disease gets worse. However, you may:

- have less energy
- have trouble thinking clearly
- have a poor appetite
- have trouble sleeping
- have dry, itchy skin
- have muscle cramping at night
- have swollen feet and ankles
- have puffiness around your eyes, especially in the morning
- need to urinate more often, especially at night.

What happens if my test results show I may have chronic kidney disease?

Your doctor will want to pinpoint your diagnosis and check your kidney function to help plan your treatment. The doctor may do the following:

- Calculate your Glomerular
 Filtration Rate (GFR), which is
 the best way to tell how much
 kidney function you have. You
 do not need to have another
 test to know your GFR. Your
 doctor can calculate it from
 your blood creatinine, your
 age, body size and gender.
 Your GFR tells your doctor
 your stage of kidney disease
 and helps the doctor plan
 your treatment. (See chart
 "Stages of Kidney Disease" on
 page 13.)
- Perform an ultrasound or CT scan to get a picture of your kidneys and urinary tract. This tells your doctor whether your kidneys are too large or small, whether you have a kidney stone or tumor and whether there are any problems in the structure of your kidneys and urinary tract.

Perform a kidney biopsy, which is done in some cases to check for a specific type of kidney disease, see how much kidney damage has occurred and help plan treatment. To do a biopsy, the doctor removes small pieces of kidney tissue and looks at them under a microscope.

Stages of Kidney Disease				
Stage	Description	Glomerular Filtration Rate (GFR)*		
1	Kidney damage (e.g., protein in the urine) with normal GFR	90 or above		
2	Kidney damage with mild decrease in GFR	60 to 89		
3	Moderate decrease in GFR	30 to 59		
4	Severe reduction in GFR	15 to 29		
5	Kidney failure	Less than 15		

^{*}Your GFR number tells your doctor how much kidney function you have. As chronic kidney disease progresses, your GFR number decreases.

Your doctor may also ask you to see a specialist who will consult on your case and help manage your care.



What should I do if I am at increased risk but I do not yet have chronic kidney disease?

You should visit your doctor or clinic for regular checkups and tests for CKD.

You should also ask your doctor what you can do to lower your chances of developing kidney disease. Your doctor may tell you to:

- Carefully follow prescribed treatments to control diabetes, high blood pressure or both.
- Lose excess weight by following a healthy diet and regular exercise program.
- Stop smoking if you are a smoker.

- Avoid taking large amounts of over-the-counter pain relievers.
- Make some changes in your diet, such as eating less salt and less protein.
- Limit your intake of beer, wine and liquor.

If I have chronic kidney disease, can I keep it from getting worse?

Most likely. Early detection and treatment can often slow or stop CKD. The degree to which your treatment can achieve this goal depends on:

- Your stage of CKD when you start treatment. The earlier you start, the better off you are.
- How carefully you follow your treatment plan. Learn all you can about CKD and treatment for it, and make sure to follow all the steps of your treatment faithfully.
- The cause of your kidney disease. Some kidney diseases are more difficult to control.

How is chronic kidney disease treated?

The type of treatment depends on your stage of CKD and other health problems you may have. Ask your doctor about how to:

- Control high blood pressure carefully. You may need to take one or more medications to reach your target blood pressure. Your doctor may prescribe high blood pressure medications called angiotensin converting enzyme (ACE) inhibitors or angiotensin receptor blockers (ARBs). Studies show that these medicines help protect your kidneys. You may also need to lose excess weight and limit the amount of salt in your diet to help control high blood pressure.
- Control blood sugar carefully if you have diabetes. Studies have shown that strict control of blood sugar may help slow CKD.

- Follow a special diet that controls the amount of protein you eat. Your doctor can refer you to a registered dietitian who can help plan your meals, so that you're eating the right foods and amounts to keep you at a healthy weight.
- Treat anemia. Drugs called erythropoiesis stimulating agents (also called ESAs for short) and iron supplements may be used. ESAs help your body make a hormone called erythropoietin (EPO), which your body needs to make red blood cells. Not having enough EPO is the most common cause of anemia in patients with kidney disease. Your body also needs iron to make red blood cells. Without enough iron, your ESA treatment will not work



Prevent bone and mineral disease by keeping your phosphorus level in balance. Phosphorus is an important mineral in your body. Your kidneys normally remove excess amounts of phosphorus from your blood. When your kidneys are not working properly, high levels of phosphorus can build up in your blood stream. This leads to calcium loss, which weakens your bones and allows them to break easier. It can also lead to heart disease. To help prevent this,

your doctor may tell you to limit the amount of high-phosphorus foods you eat, take a type of medication called a phosphate binder with your meals and snacks, and take a special prescription form of vitamin D.

- Follow an exercise program approved by your doctor.
- Take steps to prevent heart problems. This may include treating diabetes, high blood pressure and anemia, and lowering your cholesterol level if it is too high.
- Stop smoking if you are a smoker. Smoking makes heart and kidney disease worse.
- Visit your doctor or clinic regularly. This should include checking your GFR, urine protein and nutritional health. Ask your doctor about your results and keep track of them.

What happens if kidney failure occurs?

If CKD gets worse and kidney failure occurs, two successful treatments can be donedialysis or a kidney transplant. These treatments can help you stay healthy and continue your daily activities. Dialysis is a type of treatment that removes wastes and excess fluid from your blood. There are two different types of dialysis you can choose from hemodialysis and peritoneal dialysis. A kidney transplant is an operation to place a new kidney in your body to take over the work of your failed kidneys. The kidney may come from someone who has died or from a living donor who may be a close relative, friend or possibly a stranger who wished to donate a kidney to anyone in need of a transplant. For more information about these treatments, see the National Kidney Foundation resources on pages 24-25.



Is there anything else I should know?

Studies show that if you have persistent protein in your urine, you also have an increased chance of having heart problems. You should speak to your doctor about how to reduce your risk of heart disease. This may include steps such as:

- controlling diabetes and high blood pressure
- treating anemia
- lowering a high cholesterol level. (To do this, you may need to follow a diet and exercise program and possibly take medications to lower cholesterol.)

Depending on your symptoms, your doctor may order more tests to check on your heart.

Key points to remember:

- Some things can increase your risk for CKD. If you have any of the risk factors listed on page 8, make sure to visit your doctor or clinic and get tested. Early detection and treatment can keep CKD from getting worse.
- People who are at increased risk for CKD should have their blood pressure checked. They should also have a simple test for protein in the urine and a blood test for creatinine.
- Persistent protein in the urine (two positive tests over several weeks) is an early sign of CKD. Your doctor will want to pinpoint your diagnosis and check your kidney function to help plan your treatment.

- The best way to check your kidney function is to estimate your GFR (glomerular filtration rate). You do not need another test to do this. Your doctor can calculate your GFR from the results of your blood creatinine test and your age, body size and gender.
- If you are at increased risk, but do not have CKD, you should ask your doctor how to reduce your chances of developing kidney disease.
- If you have persistent protein in your urine, you may also have an increased risk for heart disease. Speak to your doctor about how to lower your risk for heart disease.
- Contact your local NKF affiliate, or call the national toll-free number (800.622.9010) to find out if Kidney Early Evaluation Program (KEEP®) is available in your area.

What other information is available?



If you have questions, speak with your healthcare team. They know you and can answer questions about you.

If you want to read more about kidney disease, the National Kidney Foundation has more than 50 other publications that cover many subjects, such as:

- CKD risk factors like hypertension and diabetes
- Complications of chronic kidney disease, such as cardiovascular disease, anemia or bone problems

- Nutrition for CKD patients, with information about carbohydrates, protein, sodium, phosphorus and potassium
- Treating kidney disease early
- Treating kidney failure with transplantation or dialysis.

There are two ways to learn about the many free resources available to you:

- Call the National Kidney Foundation at 800.622.9010.
- Visit the National Kidney
 Foundation website
 (www.kidney.org/store).
 (All publications are free,
 but there is a limit of five per person.)

Becoming an educated patient is very important to being healthy!

KNOW YOUR GFR AND OTHER IMPORTANT NUMBERS

TEST	Results/ Date	Results/ Date	Results/ Date
GFR			
Urine protein			
Blood pressure			
Blood sugar (glucose)			
Other			

NOTES


National Kidney Foundation

The National Kidney Foundation (NKF) is dedicated to preventing kidney diseases, improving the health and well-being of individuals and families affected by these diseases and increasing the availability of all organs for transplantation.

With local offices nationwide, the NKF provides early detection screening and other vital patient and community services. The Foundation conducts extensive public and professional education, advocates for patients through legislative action, promotes organ donation and supports kidney research to identify new treatments.

In 2009 NKF launched a groundbreaking multifaceted collaborative initiative to "END THE WAIT!" for a kidney transplant in the United States in 10 years by using proven strategies to eliminate barriers to donation and institute best practices across the country.

The NKF relies on individual and corporate donations, foundation and government grants, membership and special events to support its range of programs, services and initiatives.







SIEMENS HEALTHCARE DIAGNOSTICS



National Kidney Foundation 30 East 33rd Street New York, NY 10016 800.622.9010

www.kidney.org



We proudly state that this publication is printed on recycled paper that contains 10% post-consumer waste. The paper is also Forest Stewardship Council (SFC) certified which guarantees it is made from trees that come from carefully managed and responsibly harvested forests that adhere to strict environmental and socioeconomic standards. Please recycle this publication when you are finished with it.